



“杏輝”

PIC/S GMP 藥品

貝他每麗 軟膏

BETASAY Ointment “Sinphar”

—— 外用消炎劑 ——
(Anti-inflammatory Agent)

Betamethasone dipropionate 是一種合成的皮質類固醇，因為具消炎、止癢和血管收縮的作用，因此可有效治療各類發炎性皮膚病。軟膏內添加 Salicylic acid 具有去角質作用。

成分：每公克含：

Betamethasone Dipropionate.....0.64mg
(eq. to Betamethasone..... 0.50mg)
Salicylic Acid.....30mg

賦形劑：

White Petrolatum、Liquid Paraffin、PEG-20 Stearate、Beeswax、Propylene Glycol、Butylated Hydroxytoluene、Butylated Hydroxyanisole.

適應症：

慢性異位性皮膚炎、神經性皮膚炎(單純苔癬)、牛皮癬、濕疹、扁平苔癬、頭皮皮脂漏皮膚炎、汗皰、尋常性魚鱗癬等發炎症狀之解除。

用法用量：

- 1.本品應每天使用1~2次，以適量均勻塗抹於患處。
- 2.每星期使用量不應超過60gm。

副作用：

灼熱感、癢、皮膚刺激、乾燥、毛囊炎、多毛、起疹及低色素沈著等不適反應很少發生。

注意事項：

- 1.長期使用皮質類固醇製劑，將造成皮膚萎縮或產生條紋，若有此情況發生，請立即停藥。
- 2.若有皮膚刺激、過敏、過乾或皮膚剝落情形發生，請立即停藥。
- 3.使用本品於大面積損傷皮膚，將產生皮質類固醇全身性吸收的危險。
- 4.一旦發生感染，應給予適當之抗菌藥物治療。
- 5.若症狀於使用本品一週後未見改善，請立即停藥。
- 6.孕婦使用的安全性尚未建立，除非絕對必要，孕婦應避免大量或長期使用本品。
- 7.避免軟膏靠近或直接接觸到眼部。

禁忌：

- 1.本品不適用於牛痘、水痘、單純性疱疹等病毒感染及黴菌感染之皮膚疾患。
- 2.已知對 Betamethasone dipropionate 及 Salicylic acid 成份過敏者禁用。

貯藏：室溫下保存，避免日光直射及受潮，並置放於兒童不易取得之處。

本藥須由醫師處方使用

衛署藥製字第042269號
G-6310號

包裝：1000公克以下鋁管、塑膠罐裝。



杏輝藥品工業股份有限公司

SINPHAR PHARMACEUTICAL CO., LTD.

台灣宜蘭縣冬山鄉中山村中山路84號

84, CHUNG SHAN ROAD, CHUNG SHAN VILLAGE, TUNG - SHAN SHINE, I - LAN, TAIWAN

服務專線：(0800)015151

website : <http://www.sinphar.com>

BETASAY Ointment "Sinphar"

Anti-inflammatory Agent

Betamethasone dipropionate, a synthetic corticosteroid, is effective in the treatment of corticosteroid responsive dermatoses because of their antiinflammatory, anti-pruritic, and vasoconstrictive actions. The addition of salicylic acid exerts a keratolytic on the epidermis.

Composition : Each gm Contains :

Betamethasone Dipropionate.....0.64mg
(eq. to Betamethasone..... 0.50mg)
Salicylic Acid.....30mg

Other ingredients :

White Petrolatum · Liquid Paraffin · PEG-20 Stearate · Beeswax ·
Propylene Glycol · Butylated Hydroxytoluene · Butylated
Hydroxyanisole.

Indications :

Chronic eczematous dermatitis, neurodermatitis, psoriasis, eczema,
lichen planus, seborrheic dermatitis of scalp, pompholyx,
ichthyosis vulgaris.

Dosage and Administration :

1. Apply thinly 1~2 times daily.
2. Max. 60gm per week.

Side effect :

Burning, itching, irritation, dryness, folliculitis, hypertrichosis,
acneiform eruptions and hypopigmentation may occur rarely.

Precautions :

1. Prolonged use of corticosteroid preparations may produce striae or atrophy of the skin or subcutaneous tissue. If this occurs, treatment should be discontinued.
2. If irritation, sensitization, excessive dryness, or unwanted scaling develop treatment should be discontinued.
3. Application over extensive lesions may result in significant systemic absorption producing hypercortisonism manifesting.
4. If an overt infection is present, appropriate antimicrobial treatment is indicated.
5. If symptomatic response is not noted within a few days to a week, the local application should be discontinued.
6. The safety of topical corticosteroid use in pregnant women has not been established, drugs of this class should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. Drugs of this class should not be used extensively in large amounts or for prolonged periods of time in pregnant patients.
7. It should not be used in or near the eyes.

Contraindications :

1. It should not be used in viral diseases including vaccinia, varicella, herpes simplex, and fungal infections; also tuberculosis of the skin.
2. Hypersensitivity to any one of the components.

Storage :

Store at room temperature. Keep all drugs out of reach of children.



SINPHAR PHARMACEUTICAL CO., LTD.

84, CHUNG SHAN VILLAGE, TUNG - SHAN SHINE, I - LAN, TAIWAN
web site : <http://www.sinphar.com>